




MDL Patent Chemistry Database

Positioning, Content, Competitive Advantages
[Confidential]

Presented by: Eva Seip
Title: Senior Product Manager
Date: 27-October-2004




MDL® Patent Chemistry Database

What is it?

- A new structure-searchable CrossFire database
 - mainly for end-users, but with interesting features for information professionals also
 - indexing chemical reactions, substances and substance related information from Chemistry and LifeScience patents (World, U.S. and European) since 1976
 - addressing the following business critical issues:
 - **More effective synthesis planning**
 - **Better bioactivity profiling**
 - **Easy patent relevance check**
- Integrated with other data sources on DiscoveryGate soon (mid 2005)





Content and Coverage

- **Covered International Patent Classes [IPC]**

C07 Organic Chemistry*

A61K Drugs [Medicinal, Dental, Cosmetic Preparations]

A01N Biocides [Agrochemicals, Disinfectants, etc.]


C09B Dyes [can be pharmacological active]

* Polymers indexed from 2004 onwards, when mentioned in these 4 IPC's (e. g. formulations)


- **Covered Patent Agencies**


- World (WO), United States (US), European (EP) patents from Publication Year 1976 onwards





Database Design – Three “Contexts”





Competitive Advantages - Citations

Database Content:

At release: approx. 340,000 Patent Citations

Anticipated Growth: 35,000 Patents / Year

Your Benefits: Easy Relevance Check

- **Full claims text** together with Main Markush structures and main Markush Reactions*
- **Expanded Markush Structure** with full substituent list
- Defined **compounds related to Markush** structure
- Easy relevance check without time consuming access to original patent document

* On release, available for patents published since December 2003





Claim & Markush Structures to explore fast scope of patent

Title / Abstract / Claims

Title 1-substituted 2,5-dithienyl pyrrole derivatives and film-forming materials

Abstract A 1-substituted 2,5-dithienylpyrrole derivative having the following formula (I), in which R is hydrogen, a substituted or non-substituted alkyl group, or a substituted or non-substituted aromatic group, Y is hydrogen or cyano group, it can be involved the case that one of Ys may be hydrogen and the other may be cyano group, and n is an integer of 1 to 3. The derivative is used for forming films.

Claims

What is claimed is:

1. A 1-substituted 2,5-dithienylpyrrole derivative having the following formula (I).

[Figure]

in which R, is hydrogen, a substituted or non-substituted alkyl group, or a substituted or non-substituted aromatic group, Y is hydrogen or cyano group, provided that one of Ys may be hydrogen and the other may be cyano group, and n is an integer of 1 to 3.


Claim Text

Language English
Number of pages 8
Manually excerpted yes


Markush Structures

Markush PRN [19](#), [48](#), [49](#), [50](#)


PRN=19



PRN=48



PRN=49



Markush*
Structure
Display

* On release, available for patents published since December 2003



Highlight: Expanded Markush Structure

Substance Characterization

Patent Compound Registry Number **305**


Substance Type **Markush**
organic compound

Entry Date (YYYY/MM/DD) 2003/10/16

Update Date (YYYY/MM/DD) 2003/10/16

Referencing Compounds [click here](#)


PRN=19



Compressed MARKUSH: Click to expand details

Markush Viewer

To Report Print Help OK



Label	Value	Size	Attributes	Substitution	Frequency
Y	H				
	CN				
R	H				
	alkyl	1-30C	os		
	aromatic group		os		
	formyl				
	acyl				
	alkoxycarbonyl				
n	alkenyl				
	1-3				

Generic Symbols as written in Full-text



* On release, available for patents published since December 2003



Defined Substances related to Markush - Easy relevance check

Substance Characterization

Patent Compound Registry Number **305**

Substance Type **Markush**


organic compound

Entry Date (YYYY/MM/DD) 2003/10/16

Update Date (YYYY/MM/DD) 2003/10/16

Referencing Compounds **[click here](#)**


PRN=19




Compressed MARKUSH: Click to expand details

Representatives of the given Markush Structure:


Hit 1 PRN=9 C70H70NeS4




Hit 2 PRN=11 C28H14NeOS2




Hit 3 PRN=13 C28H10FN7S2



Hit 4 PRN=14 C28H10ClN7S2



* On release, available for patents published since December 2003



Competitive Advantages - Reactions

Approx. 1.5 M reactions (structure-searchable; 1976 onwards)

Strong growth anticipated: ~ 500,000 reactions/year


Your Benefit: More Effective Synthesis Planning

- **Complete Reaction Descriptions**
Immediately reproduce chemistry without the need to order the patent document
- **InfoChem ClassCodes**
Link to Similar Reactions in all other MDL Databases
- **Markush Reaction Display***
Explore the scope of patent coverage and reaction scope



* On release, available for patents published since December 2003

Experimental Text



Reaction Details

Reaction Classification

Preparation

Example Text

EXAMPLE 1

methyl 4-[2-[4-(4-pyridyl)piperazin-1-yl]acetyl]phenoxyacetate

A solution of methyl 4-bromoacetylphenoxyacetate (4.3 g) in acetonitrile (50 ml) was added dropwise over 40 minutes to a stirred solution of 1-(4-pyridyl)piperazine (4.9 g) in acetonitrile (100 ml).

Stirring was continued for a further 1.5 hours, then the solution was filtered and the filtrate evaporated in vacuo.

The solid residue was triturated with water (50 ml), then dried and suspended in methylene chloride (50 ml).

The suspension was then filtered and the filtrate concentrated to a small volume.

Purification by flash chromatography on neutral alumina eluting first with dichloromethane, then 0.

5percent w/v methanol/dichloromethane and finally 1 percent w/v methanol/dichloromethane gave the title

compound, 1.93 g, as a solid: m.p. 150.deg.-152.deg. C.; NMR(d6DMSO) δ 8.14(2H,d), 7.98(2H,d), 7.03(2H,d), 6.78(2H,d), 4.90(2H,s), 3.83(2H,s), 3.72(3H,s), 3.34(4H,bt), 2.65(4H,bt); m/e 370 (M+H)+; calculated for C₂₀H₂₃N₃O₄: C, 65.0; H, 6.3; N, 11.4. found: C, 65.2; H, 6.4; N, 11.3percent.


Easy relevance check

- Example Number
- Example Title
- Example Text
- Spectral Data (Product)

Find Similar Reactions in (other) db's

Reaction Identification

Reaction RN	4966189
Reactant PRN	3259400 2-(3-Chloro-4-methanesulfonylphenyl)-N-(5-(tetrahydro-pyran-4-yl)-propionamide
Product PRN	3412781 2-(3-chloro-4-methanesulfonylphenyl)-N-[5-(N-pyrazin-2-yl)-3-(tetrahydropyran-4-yl)propionamide
Reaction Specification	full reaction
Entry Date (YYYY/MM/DD)	2004/03/28
Update Date (YYYY/MM/DD)	2004/03/28
Find Similar Reactions	click here




Find similar reactions in:

Select product and similarity level:

Rxn Similarity	Broad	Medium	Narrow	All Choices
compares...	reaction sites only	reaction sites and alpha atoms	reaction sites and alpha and beta atoms	search using all 3 similarities
Product 1	broad	medium	narrow	find all

5.21.2.4.1 From nitriles


Nitriles (**293**) are the most frequently employed starting materials for this class of compounds. They are typically prepared from nitriles with sodium carbonate, sodium or potassium hydroxide, or sodium ethoxide to give (**294**) (Equation (85)) [8]. To avoid the separation of the product from sodium hydroxide, anhydrous methanol or ethanol is used [130]. The highest yields are obtained when 15% excess of hydroxylamine in butanol is used and the mixture is left for 48 h at 60 °C. The product separates as a practically pure crystalline material [189]. Diamidoximes (**297**) can, in principle, be prepared according to the same method from dinitriles, such as cyanogen (**295**) with hydroxylamine or alternatively, from its addition compounds with aniline, for example, diphenyloxamine (**296**), which is treated with hydroxylamine hydrochloride (Scheme 34) [16] [17][26]. The best yields and the purest products are obtained if gaseous cyanogen is led directly in an aqueous hydroxylamine solution at 0 °C [62CVR155]. Polyoximidic amides [172] and polymers containing the oximidic amide function have been prepared from polyacrylonitrile of low molecular weight with a slight excess of hydroxylamine [158] [164].



Markush Reaction Display

Hitset Family Go [icon] [icon]

Reaction Identification	
Reaction RN	16
Reactant PRN	286
	367
Product PRN	285
Reaction Specification	Markush Reaction
Entry Date (YYYY/MM/DD)	2003/10/16
Update Date (YYYY/MM/DD)	2003/10/16
Find Similar Reactions	not available




Reaction Details	
Topic of Interest	Preparation
Example Name	Scheme B
Location in Patent	Page 7
Product PRN	285
Stage Number	1
Reactant PRN	286
	367

Ref. 1	Frontpage/Claim: 13 , Fulltext: LitLink ; Patent; Publ.: EP1174421 A1 (2002/01/23), Appl.: EP00911405.9 (20000328)
--------	--

Markush* Reaction Display

On release for patents published December 2003 -

Location in Patent
"Page Number"




Competitive Advantages - Compounds

- **1.6 million compounds** with data since 1976
- **Strong growth anticipated:** ~ 800,000 Compounds/Year

Your Benefit: Better Bioactivity Profiling

- **Numerical bioactivity data**
- Indexing of compounds not covered elsewhere:
Indexing of Prophetic Compounds since 1976
 - could be made analogously to given methods
 - have a structure, but no data associated
 - normally only accessible via Markush Structure Search
 - 20-25 % more compounds than in other databases





Competitive Advantages - Compounds

Prophetic Compounds

Synthesis of methyl-(S)-2-(2-naphthalenesulfonylamino)-3-(4-hydroxyphenyl)propionate

[0029] Methyltyrosinate(1g, 4.32mmol) was suspended in 5ml of methylene chloride, pyridine(1.4ml, 17.28mmol) was added thereto, and the reaction mixture was allowed to stand until the mixture became thoroughly transparent. After the reaction solution became transparent, trimethylsilylchloride(1.1ml, 8.64mmol) was slowly added thereto at room temperature. After 1 hour, 2-naphthalenesulfonylchloride(1.08g, 4.75mmol) was added and the resulting mixture was stirred for 24 hours. To this mixture was added 3N HCl solution, which was then stirred for about 2 hours. The organic layer was separated, the aqueous layer was extracted with methylene chloride, and the organic layers were combined. The combined organic layer was dried over Na₂SO₄ and the organic solvent was removed by distillation under reduced pressure. The resulting residue was recrystallized from toluene to give 1.29g(Yield 79%) of the title compound as a white solid.

Compounds with data (e.g. yield, spectra)

¹H NMR(CDCl₃, 300MHz) δ 8.33(s, 1H), 7.91(m, 3H), 7.68(m, 3H), 6.90(d, J=8.4Hz, 2H), 6.62(d, J=8.4Hz, 2H), 5.18(d, J=9.1Hz, 1H), 4.92(br.s, 1H), 4.21(dt, J=15, 6.0Hz, 1H), 3.37(s, 3H), 3.00(dd, J=13.5, 5.7Hz, 1H), 2.93(dd, J=14.3, 6.5Hz, 1H).


The following compounds could be prepared according to the similar procedure as Preparation 2 above

Methyl-(S)-2-(*p*-toluenesulfonylamino)-3-(4-hydroxyphenyl)propionate.
Methyl-(S)-2-(benzenesulfonylamino)-3-(4-hydroxyphenyl)propionate.
Methyl-(S)-2-(benzothiazole-2-sulfonylamino)-3-(4-hydroxyphenyl)propionate,

**Prophetic
Compounds**

Numerical Bioactivity Data

Bioactivity Data 1 of 2	
Effect	inhibition of human P38 kinase alpha
Class of Effect	Pharmacology
Type	IC50
Value of Type (mole conc.-unit)	1.98 µmol/l
Species (Scientific Name)	PHAS-I (phosphorylated heat and acid stable protein kinase inducible)
Named Method	In Vitro Assay: PHAS-I
Method Details	96 well plates; biotinylated PHAS-I was used as substrate (c=1.5 µM); activated human p38 kinase alpha (c=0.3 µM); gamma 32P-ATP (activity 1.2 µCi per 50 µl); title comp. prediluted in DMSO (1percent final conc.); incubation either for 1 hour or overnight at 30 degC; capture of biotinylated PHAS-I with 32P incorporated using high capacity streptavidin coated filter place; scintillation detection.
Location in Patent	Table 1
Ref. 1	Frontpage/Claim: 192 , Fulltext: LilLink ; G.D. Searle and Company; Publ.: US6335336 B1 (2002/01/01), Appl.: US09561423 (2000/04/28)






**Location in Patent
"Page Number"**

* From Publication Year 2004 onwards




Export to Structure-Activity-Table (SAR)

Structure	Compound RegNo	Molecular Formula	Class of Effect	Effect	Type	Value (µMol/l)	Species (Scientific Name)	Location in Patent	Citation
	179361	C22H21ClN2O3	Pharmacology	antiviral	IC50	1.8	HCMV polymerase	Page column 7-8	253; Patent; Pharmacia and Upjohn Company; Publ.: US6340680; B1; (2002/01/22), Appl.: US2000-672472; (2000/09/28)
	179360	C22H21ClN2O4	Pharmacology	antiviral	IC50	0.31	HCMV polymerase	Page column 7-8	253; Patent; Pharmacia and Upjohn Company; Publ.: US6340680; B1; (2002/01/22), Appl.: US2000-672472; (2000/09/28)
	179359	C22H17ClN2O4	Pharmacology	antiviral	IC50	1.2	HCMV polymerase	Page column 7-8	253; Patent; Pharmacia and Upjohn Company; Publ.: US6340680; B1; (2002/01/22), Appl.: US2000-672472; (2000/09/28)
	179351	C24H24ClN3O4	Pharmacology	antiviral	IC50	0.48	HCMV polymerase	Page column 7-8	253; Patent; Pharmacia and Upjohn Company; Publ.: US6340680; B1; (2002/01/22), Appl.: US2000-672472; (2000/09/28)


Predefined Export Forms

Substance Profile



NMR	
Nucleus	1H
Signals given	yes
Solvent PRN	502 chloroform-d1
Location in Patent	Page 231; 236
Original String	H-NMR (ppm, [CDC13]): 7.69 (1H, d, 4.0 Hz), 7.08 (1H, s), 6.99 (1H, d, 6.0 Hz), 4.08 (1H, d, 8.0 Hz), 3.84 (1H, t, 8.0 Hz), 3.73 (1H, d, 8.0 Hz), 3.25 (1H, t, 8.0 Hz), 1.02 (9H, s), 0.72 (6H, t, 7.0 Hz).
Ref. 1	Frontpage/Claim: 240650 , Fulltext: LitLink ; Patent; ELI LILLY AND COMPANY; Publ.: WO2003/101978 A1 (2003/12/11), Appl.: WO2003-US14539 (2003/05/22)
Bioactivity Data 1 of 2	
Class of Effect	Pharmacology
Effect	osteocalcin activation
Type	EC50
Value of Type (mole conc.-unit)	0.07535 μ mol
Cell Line / Test System	osteoblast-like cell line RG-15 (ROS 17/2.8)
Method Name	OCN (osteocalcin) Promoter Assay
Method Details	rat osteoblast-like cell line; cell were trypsinized (0.25percent trypsin) and plated into white opaque 96-well plates; after 24 h cells treated with title comp., after 48 h treatment cells were lysed and assayed for luciferase activity using Luciferase Reporter Gene Assay kit
Location in Patent	Page 353-364; 368-376
Ref. 1	Frontpage/Claim: 240650 , Fulltext: LitLink ; Patent; ELI LILLY AND COMPANY; Publ.: WO2003/101978 A1 (2003/12/11), Appl.: WO2003-US14539 (2003/05/22)
Application Data	
Area of Use	Pharmaceuticals
Use	Drug acting on osteoporosis useful for treating or preventing disease states responsive to Vitamin D receptor ligands; displays the desirable cell differentiation and antiproliferative effects of Vitamin D receptor ligand with reduced calcium mobilization (calcemic) effects
Preferred Administration Form	Emulsion, Ointment, Tablet, Capsule, Pill, Powder, Lozenge, Syrup, Aerosol
Preferred Route of Application	oral
Preferred Dosage (nonnumeric)	0.0001 - 50 mg/(kg*d)
Formulation Given	yes

A substance record is a substance profile accumulating multiple properties and reactions (from different patents)



Technical Details

- **CrossFire format database for use with**
 - CrossFire Direct
 - in-house CrossFire Server [delivery via tape, FTP]
 - DiscoveryGate (coming soon)
- **DiscoveryGate implementation**
 - DatabaseBrowser (coming soon)
 - Compound Index (coming soon)
- **Updates / Timeliness**
 - Bi-weekly updates on CrossFire Direct and DatabaseBrowser
 - Semi-automatically indexed patents with frontpage, claims, many reactions and physical data already 2 weeks after publication
 - Fully indexed record incl. Markush structures/reactions and bioactivity data a few weeks later
- **Strong Yearly Growth anticipated:**
 - + 500,000 reactions
 - + 800,000 compounds
 - + 35,000 patent citations






Alert Service – to stay up-to-date



Alert Profile: Morpholino (Alert hitsets see "Results")

Query Search Context: Substances

impl. free sites



Query Options:

Free sites on all atoms	Allow: mixtures
Stereo: off	Allow: related Markush
	Allow: salts
	Allow: additional rings
	Allow: isotopes
	Allow: charges
	Allow: radicals

Text Search
and Text Search: antivir* or biocid* Automatic truncation right

Alert
Conditions

Database: Patents Frequency: After each update

First run [date]: 2004-Feb-20 Last run: Hits [last run]: Show

Send results to: e.seip@mdl.com

Send a copy to: J.Tannemann@mdl.com

Comment: Morpholino Structures with antiviral activity Save Profile





Competitive Advantages - Summary

Aspect	Advantage
Reactions	More Effective Synthesis Planning <ul style="list-style-type: none">1.5 Million Structure-Searchable ReactionsStrong growth anticipated: + 500,000 reactions/yearComplete Experimental SectionInfoChem ClassCodes [Find Similar Reactions]Markush Reaction Display*Location in Patent* <p>* From Publication Year 2004 onwards</p>
Substances	Better Bioactivity Profiling <ul style="list-style-type: none">1.6 Million Structure-Searchable SubstancesStrong growth anticipated: + 800,000/YearProphetic CompoundsSubstance Profiles: Properties & reactions accumulated in one recordNumerical Bioactivity Data [Structure-Activity-Relationship-Table]*Related Markush Structure*Compound Identifier in Patent* & Location in Patent*
Citations	Easy Relevance Check <ul style="list-style-type: none">Full Claims Text – searchable together with structures/reactionsMarkush Structure/ Reaction Display* [Expanded Form: Substituent List]Defined Substances related to a Markush structure* ["Representatives"]



Competitive Advantages - Summary

Aspect	Advantage
Timeliness	<ul style="list-style-type: none">▪ Bi-weekly Update▪ Semi-automatically indexed patents with frontpage, claims, many compounds, reactions and phys. data already 2 weeks after publication▪ Fully indexed record incl. Markush structures, Markush reactions and bioactivity data a few weeks later▪ Alert Service [Export/Import Alert Profiles]
User-Interface	Better Integration <ul style="list-style-type: none">▪ Integration into DiscoveryGate (soon)▪ MDL Standard Look & Feel▪ Find Similar Reactions in all MDL databases [Reaction Class Codes]▪ Grouping & Sorting▪ Search & Hitset History▪ Easy to Use: Find Field Name; Predefined Search Forms▪ Expert Search Mode: Command Language▪ Sophisticated Export/Report Formats [List and Table Views]
Costs	Fixed User Fee [All-You-Can-Eat] <ul style="list-style-type: none">▪ Subscription; Alert included in Fixed Fee